


EVIDENCE REVIEW

Potential Role of Oral Rinses Targeting the Viral Lipid Envelope in SARS-CoV-2 Infection

Valerie B. O'Donnell ^{1,2,*}, David Thomas^{1,3}, Richard Stanton^{1,2}, Jean-Yves Maillard^{1,4}, Robert C. Murphy⁵, Simon A. Jones^{1,2}, Ian Humphreys^{1,2}, Michael J.O. Wakelam⁶, Christopher Fegan², Matt P. Wise⁷, Albert Bosch⁸, Syed A. Sattar⁹

¹Systems Immunity Research Institute, ²School of Medicine, ³School of Dentistry, ⁴School of Pharmacy and Pharmaceutical Sciences, Cardiff University, CF14 4XN, UK, ⁵Department of Pharmacology, University of Colorado Denver, Aurora, CO 80045, USA, ⁶Babraham Institute, Babraham Research Campus, Cambridge, CB22 3AT, UK, ⁷University Hospital of Wales, Cardiff, CF14 4XW, UK, ⁸Enteric Virus Laboratory, University of Barcelona, 08028 Barcelona, Spain and ⁹Faculty of Medicine, University of Ottawa, Ottawa, Ontario K1H 8M5 Canada

*Corresponding author. E-mail: o-donnellvb@cardiff.ac.uk

Abstract

Emerging studies increasingly demonstrate the importance of the throat and salivary glands as sites of virus replication and transmission in early COVID-19 disease. SARS-CoV-2 is an enveloped virus, characterized by an outer lipid membrane derived from the host cell from which it buds. While it is highly sensitive to agents that disrupt lipid biomembranes, there has been no discussion about the potential role of oral rinsing in preventing transmission. Here, we review known mechanisms of viral lipid membrane disruption by widely available dental mouthwash components that include ethanol, chlorhexidine, cetylpyridinium chloride, hydrogen peroxide, and povidone-iodine. We also assess existing formulations for their potential ability to disrupt the SARS-CoV-2 lipid envelope, based on their concentrations of these agents, and conclude that several deserve clinical evaluation. We highlight that already published research on other enveloped viruses, including coronaviruses, directly supports the idea that oral rinsing should be considered as a potential way to reduce transmission of SARS-CoV-2. Research to test this could include evaluating existing or specifically tailored new formulations in well-designed viral inactivation assays, then in clinical trials. Population-based interventions could be undertaken with available mouthwashes, with active monitoring of outcome to determine efficacy. This is an under-researched area of major clinical need.

Key words: coronavirus; lipid; envelope; oropharynx; virus; respiratory

The Viral Lipid Envelope

In common with many viruses, such as influenza and herpes simplex, coronaviruses are surrounded by a fatty layer, called a

“lipid envelope,” into which the spike glycoproteins required for infection are inserted (Figure 1). Viral envelopes are acquired at host cell membranes—some at the plasma membrane, others at internal cell membranes such as the nuclear membrane,

Submitted: 1 May 2020; Revised: 4 May 2020; Accepted: 8 May 2020

© The Author(s) 2020. Published by Oxford University Press on behalf of the American Physiological Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Cartoon Representation of the SARS-CoV-2 Glycoprotein, Embedded in the Viral Envelope, along with Membrane Disrupting Agents. Ribbon diagram was obtained from Wrapp et al.,¹ chemical structures were from PubChem (<https://pubchem.ncbi.nlm.nih.gov/>) and Nieto-Garai et al.²

endoplasmic reticulum, and Golgi complex.^{3,4} During this, viral proteins are incorporated at the expense of host cell proteins, creating the shed viral particle.⁵ Thus, for most viruses, the envelope lipids are considered to be the same as the host membranes (phospholipids, sphingolipids, and some cholesterol). Lipid composition is not the same across subcellular membranes with mammalian plasma membranes having higher cholesterol and sphingolipid content.⁶⁻¹² While the lipid makeup of the envelope of SARS-CoV-2 has not been characterized yet, coronaviruses are known to bud from the endoplasmic reticulum Golgi intermediate compartment (ERGIC), before being transported by exocytosis in cargo vesicles.^{13,14} This indicates their composition will be related to endoplasmic reticulum membrane, which contains more phosphatidylcholine, but less cholesterol and sphingolipids than the plasma membrane.⁶⁻¹² A recent report demonstrated that coronavirus

(HCoV-229E) regulates host lipid metabolism in response to infection, in common with many other viruses.¹⁵⁻¹⁷ However, no information on the virus lipid envelope composition was provided, and its specific composition has not been determined experimentally.

The Soap/Alcohol Virucidal Public Health Advice Relating to Surface Neutralization

It is widely known that interfering with the lipid envelope represents a virucidal strategy to target many coronaviruses, with a large body of work evidencing the impact of many agents.^{18,19} For a summary, refer to Kampf et al., a systematic review providing tables showing data from different original publications for inactivation of coronaviruses by biocidal agents in suspension tests.¹⁹ During the 2003 SARS-CoV outbreak, viral material

was detected on hospital surfaces, leading to the idea that surface decontamination would be an important approach. At that time, various compounds were considered, including ethanol at high concentrations of 60%–70% (v/v), since these doses had been found to be highly effective against several viral pathogens, including coronaviruses.^{19–21} Recent studies on SARS-CoV-2 also support this, with high concentrations being highly effective as shown in a recent preprint.²² The historical reason for only testing high concentrations in microbicidal research has been that these give broad-spectrum activity toward bacteria, viruses, and fungi, and thus for use on inanimate surfaces/fomites where the target microbes are unknown would be always preferred. The consensus view is that enveloped viruses, such as herpesviruses, orthomyxoviruses, paramyxoviruses, and coronaviruses are highly sensitive to 60%–70% (v/v) ethanol with almost immediate inactivation, while nonenveloped viruses are less or not susceptible.^{19–22}

We are now widely encouraged to use soap or 60%–70% alcohol-based gels to inactivate SARS-CoV-2, based on the view that these agents damage the lipid envelope, including in recent World Health Organisation (WHO) and Environmental Protection Agency recommendations (<https://apps.who.int/iris/bitstream/handle/10665/331138/WHO-WPE-GIH-2020.1-eng.pdf>; <https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2>; <https://www.who.int/gpsc/tools/GPSC-HandRub-Wash.pdf>). At the same time, there has been no discussion of oral antiviral strategies, apart from a recent response to an article in the *British Medical Journal* calling for protection for healthcare workers against infection (<https://www.bmj.com/content/369/bmj.m1324/rr-5>). Properly designed clinical trials that address this issue are currently lacking in the literature. Current WHO interim guidance on clinical management of SARS-CoV-2 in the home is focused on the use of personal protection, including face masks, along with hand, clothing, and surface sanitation, to reduce risk of airborne and direct spread of the virus, but does not mention oral hygiene (<https://apps.who.int/iris/handle/10665/331133>). Thus, its utility in the setting of SARS-CoV-2 has not been considered systematically, and there is a lack of either positive or negative robust clinical evidence.

Mouthwashes vary widely in composition; however, some commercially available formulations contain ethanol at 14%–27% (w/v) in the United Kingdom, Europe, and the United States. Prompted by this, we reviewed the available scientific literature to establish whether oral treatment using ethanol-based or other types of mouthwashes could present a strategy to either dampen or reduce viral load, to potentially restrict virus transmission in the current pandemic situation, particularly for vulnerable individuals or healthcare workers. We found that there is a paucity of data systematically testing the impact of lower (less toxic) ethanol concentrations on enveloped virus inactivation, with most simply reiterating the use of the higher concentrations described above.^{18,20,21,23–26} We also found a paucity of robust clinical studies in this area that address in a randomized double-blind manner the impact of oral rinsing on objective measures, specifically neutralization of enveloped viruses, including coronaviruses.

Viral Load, Saliva/Throat Virus, and Disease Severity in SARS-CoV-2

It is becoming increasingly recognized that the throat is a major site of replication and shedding of virus in COVID-19 illness,

and that viral load is important.²⁷ Throat and sputum are abundant in particles, which peak 5–6 days after symptom onset, and decline thereafter.^{28,29} Viral load correlates with older age,²⁹ and a study of 76 patients in Nanchang, China, showed that those with severe SARS-CoV-2 tend to have higher viral load and longer virus-shedding period than those with mild disease.³⁰ Similarly, viral load was linked with lung disease severity in a study of 12 patients with pneumonia.³¹ Many asymptomatic individuals have modest levels of detectable viral RNA in their oropharynx for at least 5 days, which is similar to individuals with clinical symptoms.³² Data from GTEX gene expression data indicate that angiotensin converting enzyme (ACE2) (a key receptor for COVID-19) expression is higher in salivary glands than lungs, suggesting that these could be a major source of new viral particles.³³ A recent study using mobility data and Bayesian inference inferred that a high rate of undocumented infections is responsible for rapid spread of SARS-CoV-2.³⁴ Taken together, these data suggest that the potential for transmission is high early in the disease. While further studies are needed to better understand the relationship between viral load and symptom severity, it is expected that higher levels of viral shedding in the throat or lungs might be associated with an increased ability to infect others. To date, the relationship between lung and throat viral load in terms of disease severity, is not clear, and how dampening throat virus load may impact on resulting lung disease or viral transmission is not known.

The route of SARS-CoV-2 infection is currently considered to be via respiratory droplets, similar to SARS-CoV,³⁵ and the virus particle is viable in aerosols for up to 3 h.³⁶ Although we do not yet know the minimal infectious dose, the high rate of transmission indicates this is likely to be relatively low. If correct, then strategies to reduce the number of infective virus particles in mucous membranes through promoting their removal or inactivation could contribute to reducing risk of transmission. Thus, assuming that the throat is a major site of replication in early stages (even before symptoms are apparent), the oral washing using agents that could damage or destroy the lipid envelope has the potential to reduce viral load in the oropharynx.

At this time, there is incomplete information on how SARS-CoV-2 moves from the throat and nose to the lungs, and this could include (1) viral shedding, (2) the aspiration of necrotic cell debris, or (3) direct infection of neighboring cells. Assuming viral shedding is involved, the oral rinses that target the viral lipid envelope represent a potential method to remove/rinse or inactivate infective particles generated in the throat. The specific intracellular replication cycle for SARS-CoV-2 in humans is not yet known. Based on nonsynchronized replication cycles that take <24 h, virus is likely to be secreted almost constantly.^{37,38} Oral agents will impact only on virus that is extracellular or actively budding. Therefore, the persistence of treatment will be important. How long mouthwash components retain an ability to interact with biomembranes in the mouth is unclear, and more research is required.

The Impact of Lower Ethanol Concentration on Biomembranes

When considering lower (nontoxic, more economical) ethanol concentrations, the literature on mammalian cells (from where the lipid envelope originates) provides a close comparator. We also reviewed studies on model membrane vesicles comprising phospholipids such as phosphatidylcholine; however, as these are protein-free, the impact of nonlipid components on ethanol

toxicity is not accounted for. Bacterial pathogens contain very different membranes in lipid and protein composition, including lipopolysaccharides and peptidoglycans, so they are not considered here. Below we summarize the literature on impact of ethanol on cell/model membranes (Table 1).

Low Concentrations of Ethanol Cause Swelling, Interdigitation, and Leakage in Model Membranes

Biophysical studies in the 1980s and 1990s compared various alcohols (ethanol, methanol, butanol, and propanol) for their ability to perturb model phospholipid membranes. Many were optimizing generation of lipid vesicles for drug delivery; however, the toxicity of most short-chain alcohols prohibits oral use. Here, we reviewed reports on the properties of ethanol on model membranes. Few studies directly investigated lysis, instead focusing on membrane fluidity, permeability, interdigitation, thickness, and other parameters. In one study, ethanol >3.4 M (20% v/v) resulted in membranes not being considered “stable”.³⁹ Ethanol addition causes rapid swelling of phosphatidylcholine vesicles from around 30 nm diameter at 0 M, up to between 80 and 110 nm diameter, at 1.1–1.5 M (6.5%–8.8% v/v).⁵⁵ *Interdigitation* refers to the process whereby the presence of short-chain alcohols enables the methyl group of the fatty acyl chains to move beyond the midplane of the bilayer, penetrating the opposite monolayer, and appears to be an event that precedes and promotes vesicle fusion and leakage.^{56,57} Several studies demonstrate that ethanol promotes interdigitation.⁵⁸ In one, ethanol at above 2 M (11.8% v/v) led to formation of *interdigitated* phospholipid sheets from small unilamellar vesicles (SUVs), which then annealed to form larger interdigitation-fusion vesicles (IFVs).⁴⁰ This means that ethanol at this concentration can deform small phospholipid vesicles leading to fusion and formation of larger structures. During this process, leakage of contents from vesicles is seen.^{40,42,55,59} Three studies compared membranes consisting of either phosphatidylcholine alone, phosphatidylcholine/phosphatidylethanolamine mixtures, or phosphatidylcholine/cholesterol mixtures, and showed that all became permeable at ethanol concentrations around 0.6–2.1 M (3.5%–12.3%, v/v).^{43,44,55} Elsewhere, ethanol at rather lower concentrations of 86 mM (0.5% v/v) caused lysis of phosphatidylcholine vesicles during repeated cycling through phase transition temperatures.⁴¹ Partitioning of ethanol into the membrane can be altered through the presence of additional biologically relevant lipid species such as cholesterol or gangliosides.^{43,60,61} This indicates that complex biological membranes may respond very differently, and not only the presence of other lipid types, but also the impact of proteins need to be taken into account. Nonetheless, it is clear that model membranes are sensitive to ethanol at concentrations far lower than the 60%–70% currently recommended for inactivation of virus on hard surfaces, and at amounts contained in widely available mouthwashes (Section 6).

Impact of Ethanol on Mammalian Cell Membranes

We also reviewed the impact of ethanol on mammalian cells *in vitro*. Of direct relevance to coronaviruses, a study on corneal epithelial cells showed that a 30-s incubation with 20% ethanol led to around 40% loss of viability, which increased to 70% loss at 40% ethanol. There was significant leakage of intracellular contents following 20% ethanol for 30 s.⁴⁸ This short incubation

also altered inflammatory responses, differentiation, and epithelial marker expression.⁴⁸ Several studies on the impact of ethanol on red blood cells were also found. Sonmez et al. showed that around 1 M (5.9% v/v) causes ~10% cell lysis, but higher amounts were not tested.⁴⁹ A variety of effects on red cells have been shown including potassium leakage and hemolysis, at moderate concentrations around 3–4 M (18%–23.5% v/v). However, incubation times of 15 min or greater were generally used.^{50–52} Last, a study on an intestinal cell line (Caco-2) showed that ethanol >5%–10% causes loss of viability, leakage of contents, and disruption of tight junctions, with a long incubation time of 60 min.⁵³ Since the membrane composition of coronaviruses is expected to match ERGIC (Section 1), these studies provide strong evidence that low ethanol will directly impact on the SARS-CoV-2 membrane also.

So far, we found only studies that tested the impact of reduced ethanol amounts on enveloped viruses. Both were conducted *in vitro*, and show positive outcomes in relation to virus denaturation.

- In 2007, Roberts and Lloyd found that 20% ethanol completely inactivated three enveloped viruses: Sindbis, herpes simplex-1, and vaccinia, *in vitro*, while having no effect on the nonenveloped poliovirus-1.⁴⁶ Inactivation was measured by inhibition of plaque-forming units in a viral infectivity assay, but direct impact on viral envelope was not determined. This study used a rather basic system, in the absence of a soil load, which is nowadays recommended under the American Society for Testing and Materials (ASTM) Committee E35 on Pesticides, Antimicrobials and Alternative Control Agents (<https://www.astm.org/Standards/E2197.htm>), or in the United Kingdom, the equivalent British Standard (BS) Norme Européenne (EN) standard (BS EN 14476:2013+A2:2019 Chemical disinfectants and antiseptics. Quantitative suspension test for the evaluation of virucidal activity in the medical area. Test method and requirements (Phase 2/ Step 1), <https://shop.bsigroup.com/ProductDetail?pid=00000000030401479>). Also, it was conducted at 22°C, rather than the more relevant 36.8°C oral temperature, where the impact of denaturation agents would be greater.
- In 2017, Siddharta et al. tested WHO recommended formulations against enveloped viruses, including coronavirus. Focusing on WHO formulation I, which contains 85% (v/v) ethanol, 0.725% (v/v) glycerol, and 0.125% (v/v) hydrogen peroxide, they measured *in vitro* infectivity in the presence of a soil load (0.5% w/v bovine serum albumin). A 30-s exposure of a dilution containing 34% (v/v) ethanol (40% of neat) completely prevented subsequent viral replication.⁴⁷

These studies indicate that relatively dilute ethanol will be highly effective against enveloped viruses. However, there is an urgent need to determine how coronaviruses are impacted by dilute alcohol under biologically relevant conditions (mucosa, mouth, etc.), and whether in combination with nontoxic, membrane disrupting agents, oral inactivation of SARS-CoV-2 could be achieved. A minimum amount of ethanol, for example, 10%–30% (v/v) would be effective, and contact time will also be an important parameter that may reduce ethanol exposure required. Ethanol impacts membrane properties of artificial lipid membranes, causing leakage of contents even in the absence of complete lysis. The ability of the virus to infect host cells could also be modified by inducing biophysical changes to the virus membrane which impact on protein function. The spike glycoprotein which is required for SARS-CoV-2 infectivity contains a transmembrane domain that is inserted into the viral envelope,¹ and

Table 1. In vitro and in vivo data supporting the effects of ethanol on biomembranes or enveloped viruses

Reference	Study type	Ethanol	Results
Ly and Longo ³⁹	Model membrane vesicles (membrane fluidity, permeability, interdigitation, thickness, etc.)	Ethanol >3.4 M (20% v:v)	Membranes not considered “stable”; interdigitation; rapid swelling of PC vesicles
Ahl et al. ⁴⁰	Formation of <i>interdigitated</i> PL sheets from SUV	Ethanol above 2 M (11.8% v:v)	Formed larger IFVs; leakage of contents of vesicles
Hunt et al. ⁴¹	Repeated cycling through transition phase of model membranes	Ethanol 86 mM (0.5% v:v)	Lysis of PC vesicles
Komatsu et al. ^{42–44}	Leakage of dye from vesicles made of PC, PE/PC, or PC/cholesterol.	0.6–2.1 M (3.5%–12.3%, v/v)	Calcein leaks out at low ethanol concentrations. Rapid swelling of vesicles.
Dennison et al. ⁴⁵	In vitro—Herpes, influenza, rotavirus, and adenovirus	26.9% ethanol (v:v) with essential oils	Enveloped viruses (herpes and influenza) were significantly impacted
IADR abstract 2010	H1N1 Influenza A pandemic strain, in vitro	21.6% ethanol, 30-s rinse	>99.99% reduction in infectivity
Roberts and Lloyd ⁴⁶	Three enveloped viruses: Sindbis, herpes simplex-1 and vaccinia, in vitro	20% (v:v) ethanol	Completely inactivated
Siddharta et al. ⁴⁷	Enveloped viruses; in vitro infectivity WHO formulation I in the presence of coronavirus.	30-s exposure of a dilution containing 34% (v:v) ethanol	Completely prevented subsequent viral replication
Oh et al. ⁴⁸	Mammalian cell membranes: Corneal epithelial cells	20% ethanol; 30-s incubation	40% loss of viability; high level of leakage of intracellular contents
Sonmez et al. ⁴⁹ Chi and Wu ⁵⁰ and Tyulina et al. ^{51,52}	Mammalian cell membranes: Red blood cells Mammalian cell membranes: Red blood cells	1M (5.9% v:v) ethanol Moderate concentrations around 3–4M (18%–23.5%).	Approximately 10% cell lysis Potassium leakage and hemolysis
Wang et al. ⁵³	Mammalian cell membranes: Intestinal cell line (Caco-2)	Ethanol >5%–10%: long incubation time of 60 min	Loss of viability, leakage of contents, and disruption of tight junctions
Meiller et al. ⁵⁴	In vivo human study	21.6% ethanol, 30-s rinse	Recoverable virions of herpes simplex types I and II to 0 post rinse; at 30 min all lower than prerinse, 11/20 remained 0
Meiller et al. ⁵⁴	In vivo human repeat study	21.6% ethanol, 30-s rinse	0 recoverable virions in 18/20 post rinse and 12/20 at 30 min; at 60 min all less than baseline
Sattar et al. (unpublished data)	Finger pads of adults; Dried inocula; human respiratory coronavirus 229E	Hand gels with 60% and 70% ethanol exposed for 20 s	Viability titer of the virus was reduced by >99.99% in both cases

Studies cited in our text are summarized above for type, ethanol amount, and outcome. They are listed in order of model membranes, followed by in vitro studies on viruses, studies on mammalian cell membranes, then in vivo studies. Ethanol concentrations were listed also, in some cases, whether v/v or w/v was used was not provided in the study. In all studies, refer to the primary literature for full information on the impact of ethanol on the membrane. PC, Phosphatidylcholine, PE, phosphatidylethanolamine.

it is well known that lipid membrane biophysical perturbations can impact on conformation and function of many transmembrane proteins in mammalian cells. In this regard, the lipid membrane of HIV-1 was recently demonstrated to stabilize viral membrane glycoproteins and regulates their sensitivity to neutralization by antibodies.⁶² Thus, lower concentrations of ethanol could alter pathogenicity without complete neutralization of viral particles. Research is required to determine the impact of ethanol or other agents on the infective activity of the SARS-CoV-2 spike protein in vivo.

Membrane Perturbation Without Lysis can Dampen Enveloped Virus Infectivity

The concept that perturbing the membrane could inactivate viruses has recently been tested in relation to membrane-

disrupting agents, and an in vitro screen of 1000 compounds identified a series of lipidomimetics that can alter the membrane and dampen infectivity of HIV-1.² Active agents included lipids related to cholesterol, sphingosine, or aliphatic lipids with long-chain fatty acids which blocked at the stage of entry into the host cell. The impact appeared to result from the lipids being incorporated into the membrane and inducing changes in lipid order and buoyant density of the particles. In support of this, a study in the 1970s showed that fatty acids and monoglycerides of 16–18 carbon chain length are highly effective in vitro, reducing survival of herpes simplex virus to around 50% at concentrations down to 0.2 μ M.⁶³ These studies use compounds that are nontoxic to mammalian cells and show great promise, thus research is needed to determine whether they are also active against the envelope of SARS-CoV-2 both in vitro and in vivo.

Mouthwash Preparations that Show Activity Against Enveloped Viruses in Published Studies

We investigated the potential for commercially available mouthwashes to disrupt viral lipid envelopes, either due to ethanol (Table 1), or other active agents, through reviewing available literature.

Three industry-sponsored studies from the Universities of Maryland, Texas–Houston Health Sciences Centre, and State University New York tested this using a widely available formulation that combines 21%–26% ethanol with essential oils (eucalyptol 0.092%, menthol 0.042%, methyl salicylate 0.060%, and thymol 0.064% w/v). Notably, there is published evidence that eucalyptus oil and thymol have significant antiviral properties toward herpes simplex virus at these concentrations, hypothesized to relate to disruption of the viral lipid envelope.⁶⁴

- The virucidal actions of 21% (v/v) ethanol with essential oils toward an enveloped virus were reported in humans *in vivo* in 2005. A 30-s rinse reduced infectious virions of herpes simplex types I and II to effectively zero.⁵⁴ Specifically, 18/20 people demonstrated no virions post-rinse, and after a 30-min rinse, virions remained at zero for 11/20 subjects, with all subjects remaining lower than pre-rinse levels. In contrast, rinsing with distilled water reduced mean virions considerably less post 30-s rinse, and levels had largely returned to baseline by 30 min. This indicates that the mouthwash had a specific and significant impact on virion recovery. In a repeat trial, 18/20 subjects had zero virions post 30-s rinse, with 12/20 remaining at zero at 30 min. At 60 min, all 20 were still shedding virions at 1–2 log₁₀ lower than baseline, demonstrating a modest impact on viral titer. Longer contact times (eg, 60-s rinses) were not tested.⁵⁴ Herpesviruses differ from coronaviruses in that the former can erupt periodically from where they reside in the nerves; so, using mouthwash may temporarily reduce the level and it may then help promote resolution of the lesion. On the contrary, coronaviruses will be shed almost constantly when actively replicating.
- A study in 1995 tested 26.9% ethanol with essential oils against herpes, influenza, rotavirus, and adenovirus *in vitro*. Here, an impact on the viral lipid envelope was speculated since herpes and influenza were significantly impacted, while adenovirus and rotavirus (nonenveloped) were not.⁴⁵
- A follow-up unpublished study in 2010 determined that a 30-s *in vitro* exposure to 21.6% ethanol with essential oils led to >99.99% reduction of infectivity of H1N1 Influenza A pandemic strain (<https://iadr.abstractarchives.com/abstract/2010dc-131191/evaluation-of-h1n1-antiviral-properties-of-an-essential-oil-containing-mouthrinse>).

These studies provide proof-of-concept that mouthwashes containing essential oils with 21%–27% ethanol can inactivate enveloped viruses, both in the lab and in humans, with the likely mechanism being damage to the lipid envelope. Here, ethanol in combination with essential oils may provide a more effective formulation. Thus, these types of mouthwash may be effective against SARS-CoV-2, although studies have not been conducted. While other commercially available ethanol mouthwashes generally contain lower levels without essential oils, an impact on membrane biology may remain theoretically possible, and studies are required.

Chlorhexidine

Chlorhexidine is widely used for oral health in the United Kingdom, being especially effective against Gram-positive

bacteria, but to a lesser extent Gram-negative bacteria and fungi.⁶⁵ Due to its positive charge, it reacts with the negatively charged microbial surface, penetrating into the cell and causing leakage. A report on its *in vitro* virucidal effectiveness at 0.12% has indicated it can reduce the viral concentration of enveloped but not nonenveloped viruses.⁶⁶ However, this limited *in vitro* study only considered the immediate postexposure, and no further time points were included in the experimental design. Chlorhexidine is often formulated with ethanol at lower concentrations, which may in part explain its virucidal impact. A recent review of coronavirus literature identified that chlorhexidine exposure for 10 min only weakly inactivated coronavirus strains in suspension tests although the concentration used was low at 0.02%.^{19,24} Chlorhexidine formulations have been shown to retain oral antimicrobial activity for up to 12 h.⁶⁷ It is a more effective antimicrobial *in vivo* because it binds to clean oral surfaces and is released over time (substantivity).⁶⁷ Despite lower activity toward coronaviruses, a combination of chlorhexidine with alcohol may offer a useful strategy for reducing viral load over longer times.

Chlorhexidine mouthwashes have been a critical clinical tool for over 40 years to reduce oral bacterial flora and prevent infection and mucositis in cancer patients receiving chemotherapy and radiotherapy.^{68–70} However, there are no reported studies assessing the impact of mouthwashes in specifically preventing or treating viral infections in neutropenic patients. Last, a recent meta-analysis showed that chlorhexidine (rinse or gel) can reduce risk of ventilator-associated pneumonia in patients undergoing mechanical ventilation, although causative organisms were not described.⁷¹

Povidone-Iodine

Povidone-iodine (PVP-I) mouthwash has been widely studied in relation to broad-spectrum antimicrobial and virucidal actions. At 0.23%, which is routinely used in Japan, this rapidly inactivates SARS-CoV, MERS-CoV, influenza virus A (H1N1), and rotavirus *in vitro*.⁷² A second study also showed that PVP-I (0.23%) is equivalent to 70% ethanol in inactivating SARS-CoV *in vitro*.⁷³ Indeed, based on *in vitro* and limited clinical studies, in Japan, the Ministry of Health, Labour and Welfare supported daily gargling as a protective measure to prevent upper respiratory tract infections.⁷⁴ A small number of human studies supporting this in the case of PVP-I have shown reduced incidence of both bacterial and viral (influenza) infection through repeated gargling.^{72,75} In one rather limited study, the absence rate in middle schools in Yamagata City was compared over 3 months, where PVP-I gargling was encouraged in 1 school, versus 7 where it was not. A reduction of absence due to colds and influenza from a mean of 25.5% (no gargling) down to 19.8% ($P < 0.05$) was found.⁷⁶ In another study, a group of 23 patients gargled more than 4 times/day for up to 2 h. Here, acute exacerbation of chronic respiratory infection was reduced by around 50%.⁷⁵ This mouthwash is not available in the United Kingdom, although may still be purchased in Germany and other countries. As a 1% solution, PVP-I is available in Hong Kong, Korea, Singapore, Malaysia, Philippines, and Taiwan. The importance of higher concentrations of PVP-I as a broad-spectrum antimicrobial agent for topical uses is indicated by its inclusion on the World Health Organization's List of Essential Medicines (<https://www.who.int/medicines/publications/essentialmedicines/en/>). It should be noted that rare allergic reactions have been reported for PVP-I.⁷⁷

Chlorinated Water or Hypertonic Saline Rinsing

Studies from Japan surprisingly found that gargling with chlorinated tap water reduced respiratory infections, and in one, was even better than PVP-I. In one, three groups of around 130 age- and gender-matched human subjects were studied (three 15-s gargles with 20 mL, at least 3/day for 60 days).⁷⁸ Tap water reduced incidence of common cold by 36%, while PVP-I was not effective. It was speculated that chlorine in the water may have contributed, since levels in Japan were above concentrations that are known to have viricidal activity including toward enveloped species.⁷⁹ However, information on the viricidal impact of chlorine comes from in vitro studies, including 1 with a 30 min contact time, and its impact in vivo on enveloped viruses, through gargling tap water is not known.⁷⁹ In 2008, another trial calculated economics of the activity in two groups of around 120 subjects gargling water for 60 days and concluded that this was a cost-effective strategy for upper respiratory infection prevention.⁸⁰ Last, a recent study showed that gargling and nasal rinsing with hypertonic saline could reduce symptoms, duration of illness, and viral shedding. However, this was a pilot, nonblinded, self-reported study and so cannot be considered definitive.⁸¹ A follow-up study in vitro with enveloped and non-enveloped viruses including human coronavirus 229E suggested that this may have been related to altered intracellular chloride levels and peroxidase activities.⁸² None of the in vivo studies addressed the issue of which pathogens were contributing to illness, and so cannot be extrapolated to coronaviruses.

Separate to oral rinsing it is worth noting that nasal rinsing with saline is a popular method promoted to clear nasal passages for sufferers of colds and allergies. Given that virus is recovered in the nasopharynx, a similar consideration of how this might be used as preventative measure could be made. As for mouthwash, clinical studies have not systematically examined how effective nasal rinsing is for preventing respiratory infections. Notably, rare reports of serious illness when not properly cleaned, due to the presence of parasitic amoebae in unboiled tap water, has led to recommendations on careful disinfection of rinsing syringes being made by CDC (<https://www.cdc.gov/par/sites/naegleria/sinus-rinsing.html>).

Hydrogen Peroxide

Hydrogen peroxide causes oxygen-free radical-induced disruption of lipid membranes and is widely used as an agent for tooth whitening. Studies, including a recent systematic review, report that coronavirus 229E and other enveloped viruses are inactivated at concentrations around 0.5%.^{19,83} While higher concentrations of hydrogen peroxide (>5%) will induce damage to both soft and hard tissues, within the range of concentrations used in mouthwashes for whitening at 1%–3% little damage is reported.⁸⁴ Within the oral environment, hydrogen peroxide is rapidly inactivated due to the presence of host- and bacteria-derived catalase activity in saliva and other endogenous peroxidases.⁸⁵ The impact of peroxidases could theoretically be reduced by using a prerinse with water, although this is untested. A consideration with this agent is that it can have potential antiviral activities, although so far this was only seen in vitro.^{86–88}

Quaternary ammonium compounds

These are widely used as microbicidal agents that interfere with protein or lipid components on the cell surface, particularly Gram-positive or Gram-negative bacteria. Their viricidal activities are not widely reported although some reports against

enveloped viruses have been made in the literature relating to surface disinfection.⁷⁹ Among this group of compounds, cetylpyridinium chloride (CPC) was recently shown to have activity against influenza both in vitro and in vivo, through direct attack on the viral envelope, with in vitro EC₅₀ being 5–20 µg/mL.⁸⁹ CPC is used in medicated oral rinses at concentrations 0.025%–0.075% w/v (250–750 µg/mL) in the United Kingdom, while lozenges sold in some countries contain 1.4–3.0 mg of CPC.

Current Policies Relating to Oral Health and Use of Microbicides in Dental Practice and with Immunosuppressed Individuals

Dental practitioners are at elevated risk of exposure to SARS-CoV-2, and there are guidelines that advocate use of mouthwash clinically. Previously published CDC guidelines for infection control in the dental setting have cited the potential usefulness of preprocedural mouthwashes in reducing the spread of airborne pathogens of all types.⁹⁰ Indeed, studies have addressed how virucidal components including CPC and chlorhexidine can be effective in reducing bacterial contamination in this setting, although virus inactivation was not tested.^{91,92} Meng et al. in an experience-based review of their practice in Wuhan, recommended preprocedural mouthwash to reduce the oral microbial load in patients undergoing dental treatment in patients with SARS-CoV-2.⁹³ Last, two recent papers aimed at providing guidelines for endodontists in relation to SARS-CoV-2 advocated preprocedural mouth rinse with 0.2% PVP-I.^{94,95} Given that the dental community recognize, the potential for oral mouth washing in relation to reducing infection risk, extrapolating these guidelines to the wider community is worth a full discussion.

The Urgent Need for Research

Many questions need to be addressed in relation to whether oral hygiene could represent a viable approach to dampen transmission of SARS-CoV-2, and research is required to address this.

In relation to oral hygiene, we need to determine:

- Can we reduce viral load in the oropharynx through oral rinsing?
- If we can reduce load, then which oral rinse would be clinically effective: The current choice includes 20%–30% ethanol, lipid-based membrane disruptors, PVP-1, CPC, hydrogen peroxide, simple chlorinated tap water or WHO formulation I diluted to 30% of neat?
- Would a combination of agents in lower amounts be better tolerated, reducing adverse effects, and remain effective?
- What combinations or agents, contact time and frequency of use would induce antiviral activity and reduce infectivity of SARS-CoV-2?

Available research approaches include:

Statistical epidemiological studies could establish on a population level whether mouth rinsing is associated with reduced rates of throat and respiratory infections including SARS-CoV-2. Purchasing data of health-related products to model health linkages could be used. New applications to conduct wide-spread monitoring of SARS-CoV-2 symptoms, could capture use of mouthwashes to test for correlation with symptoms and severity, alongside wider purchasing sales behavior of those who are asymptomatic. Modeling approaches should

also consider population usage of mouthwash preparations and viral spread.

Underpinning research. Not all enveloped viruses are the same, and herpes, influenza, and measles viruses are considered more unstable than human coronaviruses, which may persist for up to 5 days on inanimate surfaces.^{19,36,96} Thus, research needs to focus on coronaviruses in particular. The exact composition of the SARS-CoV-2 lipidome needs to be determined using lipidomics mass spectrometry. Research should determine the impact of ethanol or other agents on the infective activity of the spike protein itself, *in vitro*, and *in vivo*. A useful virus to test *in vitro* would be the human respiratory coronavirus 229E which is used extensively as a surrogate for human coronaviruses but only requires Category (Cat2) procedures, and its replication and propagation conditions are well established already. This would be a good representative for pathogenic coronaviruses, prior to narrowing down to SARS-CoV-2 which requires Cat3 biosecurity.

The impact of temperature and soil load needs to be considered for *in vitro* studies, applying the ASTM or EU standard protocols (<https://www.astm.org/Standards/E2197.htm>). Here, the impact of not only dose/composition but also the critical issue of contact time with agents, which is a known modifiable parameter of virucidal activity can be easily tested. The virucidal mechanisms can be determined, conducting lipidomics analysis of the envelope along with assays that determine spike protein conformation and activity. While a particular ethanol concentration may achieve full inactivation, lower amounts could either help to remove virus or lead to membrane damage (permeability/leakage) that may impact throat cell virus infectivity, for example, through potential modification of the ability of the spike glycoprotein to interact with receptors on host cells. In this regard, for HIV-1, the lipid membrane stabilizes membrane glycoproteins, regulating their sensitivity to antibody neutralization.⁶² This type of action could be further enhanced if membrane disrupting agents were also included in a mouthwash. Indeed, lipidomimetic compounds have already been developed that can dampen viral infection through affecting lipid membrane structure or curvature (discussed in Section “Membrane Perturbation Without Lysis can Dampen Enveloped Virus Infectivity”).

Most virucidal research uses *in vitro* models, where the response to the agent will be different, and also does not take into account the impact of host immunity. There is an absence of animal model studies on coronavirus respiratory illness, although macaques and mice transgenic for humanized ACE2 are beginning to be studied with early indications being that both develop mild illnesses in response to SARS-CoV-2 virus (<https://www.nature.com/articles/d41586-020-00698-x>).

Clinical studies. Robustly designed, appropriately powered *in vivo* clinical studies are needed, including determination of the most effective composition. Self-reported, nonrandomized, unblinded studies are not reliable and need to be avoided. An important sequela of these over the counter medicines is that individuals may use them prior to providing diagnostic nasopharyngeal/throat swabs. This could increase the number of false-negative tests and facilitate transmission. Currently, there is no specific advice to avoid these preparations prior to testing.

Population-based interventions could be considered, although panic buying or dangerous consumption of ethanol or methanol has to be avoided. As high-risk groups come out of self-isolation, they could represent a population to evaluate clinical outcomes resulting from real-world use of available

mouthwashes. The current social restrictions will reduce a number of transmission risk factor variables and alter clinical outcomes in terms of SARS-CoV-2 infection, other respiratory infections, and adverse effects, but monitoring outcomes could provide useful data. Users could be given general advice on product use, recording timings, and duration of gargling for later analysis, or act as controls. Similarly, health workers at high risk of infection could be provided products and asked to record their use and report outcomes. Ideally, throat swabs and blood samples would be obtained for testing. There would be logistical, ethical, and regulatory issues involved in setting up investigations. However, given the theoretical plausibility and data we have reviewed plus the readily available products and urgent need to reduce SARS-CoV-2 infection, measures could be considered and action taken to instigate clinical investigation in the population during the outbreak.

Host inflammation. Mouthwashes widely utilized in daily oral and dental hygiene for cosmetic and medical reasons and have demonstrated acceptable tolerability when used multiple times daily for durations of 6 months and longer. Despite this, the impact of rinsing with these agents on throat tissue health needs to be seriously considered, since the viral lipid membrane is effectively the same as that of the host. Some of these agents, such as ethanol and hydrogen peroxide may, if used several times a day over a period of 2-3 months, induce mucosal inflammation. This was observed in a study on corneal epithelial cells, where inflammatory cytokines (IL-1b, IL-6), chemokines (IL-8/CXCL8, CCL2), and matrix metalloproteases (MMP9) were all upregulated at the mRNA levels 1-3 days after a 30-s exposure to 20% (v/v) ethanol.⁴⁸ Here, it will be important to ascertain whether a repeated daily rinse with mouthwash would have any detrimental impact on the stromal tissue lining. Alternatively, host innate immune responses in early infection could also represent a strategy to remove virus, and this has not been considered in any studies to date. Significant advances have been made into the molecular basis of alcohol-induced tissue injury. However, these studies tend to be confined to studies of acute and chronic alcohol consumption where the metabolism of alcohol into acetaldehyde and reactive oxygen intermediates modify various physiological processes linked with the maintenance of tissue homeostasis.⁹⁷ Currently, there is a lack of research into the potential impact of mouthwash on local inflammation within the throat and consideration needs to be given to both its impact on antiviral immunity and the disruption of tissue integrity.

Additional Reading

We highlight a list of excellent review articles that were consulted as part of preparing this review and were a source of primary research cited herein:

Budding of viral lipid membranes: Simons and Garoff⁹⁸

Impact of virus on lipid metabolism: Sanchez and Lagunoff⁹⁹

Review of inactivation of coronaviruses on surfaces: Wolff et al.¹⁰⁰

Emergence of SARS-CoV in 2003: Peiris et al.¹⁰¹

Two papers on the impact of ethanol on interdigitation of membranes: Slatter and Huang^{102,103}

Control of infection using Povidone-Iodine: Eggers¹⁰⁴

Summary of the composition of lipid membranes in cells: van Meer¹⁰⁵

Search strategy

Since the review covers many areas from basic biochemistry, virology, and microbicidal research, as well as clinical

information both in medicine and dentistry, multiple sources were consulted. Most references were identified from PubMed, ResearchGate, or Google, using search terms including “virus,” “coronavirus,” “lipid envelope,” “alcohol,” “membrane,” “chlorhexidine,” and others, alone and in combination. Many references that were first identified, were then investigated further to find additional source material and the original primary research which was then included. The idea for drafting the review was initiated by V.B.O. on March 21, 2020, through reaching out in person to various international experts, to get their views and input directly, via phone calls and then follow-up emails. Boots UK was approached initially to discuss the ideas and for information on formulation of available mouthwashes. Boots researchers and scientists contacted Johnson & Johnson, who provided proprietary information. Boots researchers (Kirkdale, Thornley, Povey, Inchley, and O’Shea) then conducted further searches of PubMed, Google, and ResearchGate, using the same terms. Academic and clinical expertise was consulted for virology (Stanton, Humphreys, and Bosch), clinical/ICU (Fegan and Wise), dental practice (Thomas), immunology (Jones), lipid biochemistry (Wakelem, Murphy, Simons, and O’Donnell), and microbicides (Sattar and Maillard). Phone/email correspondence with experts generated input, opinions, and identified additional references. No timeline for references was used and no languages were excluded.

Acknowledgments

The lead author gratefully acknowledges significant input from colleagues employed at Boots UK, Nottingham (Drs Andrew Inchley, Tracey Thornley, Janet Povey, Charlotte Kinchley, and Ged O’Shea), who provided formulation information on existing mouthwashes, that is not available in the public domain. Boots UK staff also linked directly with colleagues at Johnson & Johnson to obtain formulation information, and conducted literature searches, providing information to the lead author, who had full independent editorial control over drafting and editing the manuscript. The authors gratefully acknowledge expert advice on viral lipidomes from Prof. Kai Simons, Emeritus Director, Max Planck Institute of Molecular Cell Biology and Genetics, Dresden. Johnson & Johnson Ltd provided technical information on formulations. No financial agreements or other forms of payment were provided to any parties involved in preparing this review.

Funding

V.B.O. is a Royal Society Wolfson Research Merit Award Holder and acknowledges funding for LIPID MAPS from Wellcome Trust (203014/Z/16/Z).

Conflict of interest statement

None declared.

References

1. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367(6483):1260–1263.
2. Nieto-Garai JA, Glass B, Bunn C, et al. Lipidomimetic compounds act as HIV-1 entry inhibitors by altering viral membrane structure. *Front Immunol* 2018;9:1983.
3. Gerl MJ, Sampaio JL, Urban S, et al. Quantitative analysis of the lipidomes of the influenza virus envelope and MDCK cell apical membrane. *J Cell Biol* 2012;196(2):213–221.
4. Garoff H, Simons K, Dobberstein B. Assembly of the Semliki Forest virus membrane glycoproteins in the membrane of the endoplasmic reticulum in vitro. *J Mol Biol* 1978;124(4):587–600.
5. Kalvodova L, Sampaio JL, Cordo S, Ejsing CS, Shevchenko A, Simons K. The lipidomes of vesicular stomatitis virus, Semliki Forest virus, and the host plasma membrane analyzed by quantitative shotgun mass spectrometry. *J Virol* 2009;83(16):7996–8003.
6. Cluett EB, Machamer CE. The envelope of vaccinia virus reveals an unusual phospholipid in Golgi complex membranes. *J Cell Sci* 1996;109(Pt 8):2121–2131.
7. Fleischer B, Zambrano F, Fleischer S. Biochemical characterization of the Golgi complex of mammalian cells. *J Supramol Struct* 1974;2(5–6):737–750.
8. Hornick CA, Hamilton RL, Spaziani E, Enders GH, Havel RJ. Isolation and characterization of multivesicular bodies from rat hepatocytes: an organelle distinct from secretory vesicles of the Golgi apparatus. *J Cell Biol* 1985;100(5):1558–1569.
9. Howell KE, Palade GE. Hepatic Golgi fractions resolved into membrane and content subfractions. *J Cell Biol* 1982;92(3):822–832.
10. Keenan TW, Morre DJ. Phospholipid class and fatty acid composition of Golgi apparatus isolated from rat liver and comparison with other cell fractions. *Biochemistry* 1970;9(1):19–25.
11. Moreau P, Cassagne C. Phospholipid trafficking and membrane biogenesis. *Biochim Biophys Acta* 1994;1197(3):257–290.
12. Kean LS, Grant AM, Angeletti C, et al. Plasma membrane translocation of fluorescent-labeled phosphatidylethanolamine is controlled by transcription regulators, PDR1 and PDR3. *J Cell Biol* 1997;138(2):255–270.
13. Krijnse-Locker J, Ericsson M, Rottier PJ, Griffiths G. Characterization of the budding compartment of mouse hepatitis virus: evidence that transport from the RER to the Golgi complex requires only one vesicular transport step. *J Cell Biol* 1994;124(1–2):55–70.
14. Tooze J, Tooze SA. Infection of AtT20 murine pituitary tumour cells by mouse hepatitis virus strain A59: virus budding is restricted to the Golgi region. *Eur J Cell Biol* 1985;37:203–212.
15. Yan B, Chu H, Yang D, et al. Characterization of the lipidomic profile of human coronavirus-infected cells: implications for lipid metabolism remodeling upon coronavirus replication. *Viruses* 2019;11(1):73.
16. Kapadia SB, Chisari FV. Hepatitis C virus RNA replication is regulated by host geranylgeranylation and fatty acids. *Proc Natl Acad Sci USA* 2005;102(7):2561–2566.
17. Munger J, Bajad SU, Collier HA, Shenk T, Rabinowitz JD. Dynamics of the cellular metabolome during human cytomegalovirus infection. *PLoS Pathog* 2006;2(12):e132.
18. Pratelli A, Colao V. Role of the lipid rafts in the life cycle of canine coronavirus. *J Gen Virol* 2015;96(Pt 2):331–337.
19. Kampf G, Todt D, Pfaender S, Steinmann E. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect* 2020;104(3):246–251.

20. Hulkower RL, Casanova LM, Rutala WA, Weber DJ, Sobsey MD. Inactivation of surrogate coronaviruses on hard surfaces by health care germicides. *Am J Infect Control* 2011;39(5): 401–407.
21. Sattar SA, Springthorpe VS, Karim Y, Loro P. Chemical disinfection of non-porous inanimate surfaces experimentally contaminated with four human pathogenic viruses. *Epidemiol Infect* 1989;102(3):493–505.
22. Kratzel A, Todt D, V'kovski, et al. Efficient inactivation of SARS-CoV-2 by WHO-recommended hand rub formulations and alcohols. *BioRxiv*. 2020. doi: 10.1101/2020.03.10.986711.
23. Brown TT Jr. Laboratory evaluation of selected disinfectants as virucidal agents against porcine parvovirus, pseudorabies virus, and transmissible gastroenteritis virus. *Am J Vet Res* 1981;42(6):1033–1036.
24. Saknimit M, Inatsuki I, Sugiyama Y, Yagami K. Virucidal efficacy of physico-chemical treatments against coronaviruses and parvoviruses of laboratory animals. *Jikken Dobutsu* 1988; 37(3):341–345.
25. Swanson DA. A state-based regression model for estimating substate life expectancy. *Demography* 1989;26(1):161–170.
26. Wood A, Payne D. The action of three antiseptics/disinfectants against enveloped and non-enveloped viruses. *J Hosp Infect* 1998;38(4):283–295.
27. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* 2020. doi: 10.1038/s41586-020-2196-x.
28. Pan Y, Zhang D, Yang P, Poon LLM, Wang Q. Viral load of SARS-CoV-2 in clinical samples. *Lancet Infect Dis* 2020. doi: 10.1016/S1473-3099(20)30113-4.
29. To KK, Tsang OT, Leung WS, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis* 2020.
30. Liu Y, Yan LM, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* 2020. doi: 10.1016/S1473-3099(20)30196-1.
31. Liu Y, Yang Y, Zhang C, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci* 2020;63(3):364–374.
32. Zou L, Ruan F, Huang M, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382(12):1177–1179.
33. Xu J, Li Y, Gan F, Du Y, Yao Y. Salivary glands: potential reservoirs for COVID-19 asymptomatic infection. *J Dent Res* 2020;22034520918518. doi: 10.1177/0022034520918518.
34. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science* 2020.
35. Varia M, Wilson S, Sarwal S, et al. Investigation of a nosocomial outbreak of severe acute respiratory syndrome (SARS) in Toronto, Canada. *CMAJ* 2003;169(4):285–292.
36. van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med* 2020;382:1564–1567.
37. Tamin A, Queen K, Paden CR, et al. Isolation and growth characterization of novel full length and deletion mutant human MERS-CoV strains from clinical specimens collected during 2015. *J Gen Virol* 2019;100(11):1523–1529.
38. Harcourt J, Tamin A, Lu X, et al. Severe acute respiratory syndrome coronavirus 2 from patient with 2019 novel coronavirus disease, United States. *Emerg Infect Dis* 2020;26(6). doi: 10.3201/eid2606.200516
39. Ly HV, Longo ML. The influence of short-chain alcohols on interfacial tension, mechanical properties, area/molecule, and permeability of fluid lipid bilayers. *Biophys J* 2004;87(2): 1013–1033.
40. Ahl PL, Chen L, Perkins WR, et al. Interdigitation-fusion: a new method for producing lipid vesicles of high internal volume. *Biochim Biophys Acta* 1994;1195(2):237–244.
41. Hunt GR, Jones IC. A 1H-NMR investigation of the effects of ethanol and general anaesthetics on ion channels and membrane fusion using unilamellar phospholipid membranes. *Biochim Biophys Acta* 1983;736(1):1–10.
42. Komatsu H, Guy PT, Rowe ES. Effect of unilamellar vesicle size on ethanol-induced interdigitation in dipalmitoylphosphatidylcholine. *Chem Phys Lipids* 1993;65(1):11–21.
43. Komatsu H, Okada S. Effects of ethanol on permeability of phosphatidylcholine/cholesterol mixed liposomal membranes. *Chem Phys Lipids* 1997;85:67–74.
44. Komatsu H, Okada S. Increased permeability of phase-separated liposomal membranes with mixtures of ethanol-induced interdigitated and non-interdigitated structures. *Biochim Biophys Acta* 1995;1237(2):169–175.
45. Dennison DK, Meredith GM, Shillitoe EJ, Caffesse RG. The antiviral spectrum of Listerine antiseptic. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995;79(4):442–448.
46. Roberts PL, Lloyd D. Virus inactivation by protein denaturants used in affinity chromatography. *Biologicals* 2007;35(4): 343–347.
47. Siddharta A, Pfaender S, Vielle NJ, et al. Virucidal activity of World Health Organization-recommended formulations against enveloped viruses, including zika, ebola, and emerging coronaviruses. *J Infect Dis* 2017;215(6):902–906.
48. Oh JY, Yu JM, Ko JH. Analysis of ethanol effects on corneal epithelium. *Invest Ophthalmol Vis Sci* 2013;54(6):3852–3856.
49. Sonmez M, Ince HY, Yalcin O, et al. The effect of alcohols on red blood cell mechanical properties and membrane fluidity depends on their molecular size. *PLoS One* 2013;8(9):e76579.
50. Chi LM, Wu WG. Mechanism of hemolysis of red blood cell mediated by ethanol. *Biochim Biophys Acta* 1991;1062(1): 46–50.
51. Tyulina OV, Huentelman MJ, Prokopieva VD, Boldyrev AA, Johnson P. Does ethanol metabolism affect erythrocyte hemolysis? *Biochim Biophys Acta* 2000;1535(1):69–77.
52. Tyulina OV, Prokopieva VD, Dodd RD, et al. In vitro effects of ethanol, acetaldehyde and fatty acid ethyl esters on human erythrocytes. *Alcohol Alcohol* 2002;37(2):179–186.
53. Wang Y, Tong J, Chang B, Wang B, Zhang D, Wang B. Effects of alcohol on intestinal epithelial barrier permeability and expression of tight junction-associated proteins. *Mol Med Rep* 2014;9(6):2352–2356.
54. Meiller TF, Silva A, Ferreira SM, Jabra-Rizk MA, Kelley JI, DePaola LG. Efficacy of Listerine antiseptic in reducing viral contamination of saliva. *J Clin Periodontol* 2005;32(4):341–346.
55. Komatsu H, Okada S. Ethanol-induced aggregation and fusion of small phosphatidylcholine liposome: participation of interdigitated membrane formation in their processes. *Biochim Biophys Acta* 1995;1235(2):270–280.
56. Nambi P, Rowe ES, McIntosh TJ. Studies of the ethanol-induced interdigitated gel phase in phosphatidylcholines using the fluorophore 1,6-diphenyl-1,3,5-hexatriene. *Biochemistry* 1988;27(26):9175–9182.
57. Rowe ES. Thermodynamic reversibility of phase transitions. Specific effects of alcohols on phosphatidylcholines. *Biochim Biophys Acta* 1985;813(2):321–330.

58. McConnell DS, Schullery, SE. Phospholipidvesiclefusion and drug loading: temperature, solute and cholesterol effects, and, a rapid preparationfor solute-loadedvesicles. *Biochimica et Biophysica Acta* 1985;818:13–22.
59. Boni LT, Minchey SR, Perkins WR, et al. Curvature dependent induction of the interdigitated gel phase in DPPC vesicles. *Biochim Biophys Acta* 1993;1146(2):247–257.
60. Barry JA, Gawrisch K. Effects of ethanol on lipid bilayers containing cholesterol, gangliosides, and sphingomyelin. *Biochemistry* 1995;34(27):8852–8860.
61. Trandum C, Westh P, Jorgensen K, Mouritsen OG. Association of ethanol with lipid membranes containing cholesterol, sphingomyelin and ganglioside: a titration calorimetry study. *Biochim Biophys Acta* 1999;1420(1–2):179–188.
62. Salimi H, Johnson J, Flores MG, et al. The lipid membrane of HIV-1 stabilizes the viral envelope glycoproteins and modulates their sensitivity to antibody neutralization. *J Biol Chem* 2020;295(2):348–362.
63. Sands J, Auperin D, Snipes W. Extreme sensitivity of enveloped viruses, including herpes simplex, to long-chain unsaturated monoglycerides and alcohols. *Antimicrob Agents Chemother* 1979;15(1):67–73.
64. Astani A, Reichling J, Schnitzler P. Comparative study on the antiviral activity of selected monoterpenes derived from essential oils. *Phytother Res* 2010;24(5):673–679.
65. Cheung HY, Wong MM, Cheung SH, Liang LY, Lam YW, Chiu SK. Differential actions of chlorhexidine on the cell wall of *Bacillus subtilis* and *Escherichia coli*. *PLoS One* 2012;7(5):e36659.
66. Bernstein D, Schiff G, Echler G, Prince A, Feller M, Briner W. In vitro virucidal effectiveness of a 0.12%-chlorhexidine gluconate mouthrinse. *J Dent Res* 1990;69(3):874–876.
67. Bonesvold P, Lokken P, Rolla G, Paus PN. Retention of chlorhexidine in the human oral cavity after mouth rinses. *Arch Oral Biol* 1974;19(3):209–212.
68. Ferretti GA, Raybould TP, Brown AT, et al. Chlorhexidine prophylaxis for chemotherapy- and radiotherapy-induced stomatitis: a randomized double-blind trial. *Oral Surg Oral Med Oral Pathol* 1990;69(3):331–338.
69. Kuo CC, Wang RH, Wang HH, Li CH. Meta-analysis of randomized controlled trials of the efficacy of propolis mouthwash in cancer therapy-induced oral mucositis. *Support Care Cancer* 2018;26(12):4001–4009.
70. Sorensen JB, Skovsgaard T, Bork E, Damstrup L, Ingeberg S. Double-blind, placebo-controlled, randomized study of chlorhexidine prophylaxis for 5-fluorouracil-based chemotherapy-induced oral mucositis with nonblinded randomized comparison to oral cooling (cryotherapy) in gastrointestinal malignancies. *Cancer* 2008;112(7):1600–1606.
71. Veitz-Keenan A, Ferraiolo DM. Oral care with chlorhexidine seems effective for reducing the incidence of ventilator-associated pneumonia. *Evid Based Dent* 2017;18(4):113–114.
72. Eggers M, Koburger-Janssen T, Eickmann M, Zorn J. In vitro bactericidal and virucidal efficacy of povidone-iodine gargle/mouthwash against respiratory and oral tract pathogens. *Infect Dis Ther* 2018;7(2):249–259.
73. Kariwa H, Fujii N, Takashima I. Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions and chemical reagents. *Dermatology* 2006;212(Suppl 1):119–123.
74. Japan Ministry of Health LaW. Pandemic influenza preparedness action plan of the Japanese Government; 2007.
75. Nagatake T, Ahmed K, Oishi K. Prevention of respiratory infections by povidone-iodine gargle. *Dermatology* 2002; 204(Suppl 1):32–36.
76. Shiraishi T, Nakagawa Y. Evaluation of the bactericidal activity of povidone-iodine and commercially available gargle preparations. *Dermatology* 2002;204(Suppl 1):37–41.
77. Rahimi S, Lazarou G. Late-onset allergic reaction to povidone-iodine resulting in vulvar edema and urinary retention. *Obstet Gynecol* 2010;116(Suppl 2):562–564.
78. Satomura K, Kitamura T, Kawamura T, et al. Prevention of upper respiratory tract infections by gargling: a randomized trial. *Am J Prev Med* 2005;29(4):302–307.
79. Shirai J, Kanno T, Tsuchiya Y, Mitsubayashi S, Seki R. Effects of chlorine, iodine, and quaternary ammonium compound disinfectants on several exotic disease viruses. *J Vet Med Sci* 2000;62(1):85–92.
80. Sakai M, Shimbo T, Omata K, et al. Cost-effectiveness of gargling for the prevention of upper respiratory tract infections. *BMC Health Serv Res* 2008;8:258.
81. Ramalingam S, Graham C, Dove J, Morrice L, Sheikh A. A pilot, open labelled, randomised controlled trial of hypertonic saline nasal irrigation and gargling for the common cold. *Sci Rep* 2019;9(1):1015.
82. Ramalingam S, Cai B, Wong J, et al. Antiviral innate immune response in non-myeloid cells is augmented by chloride ions via an increase in intracellular hypochlorous acid levels. *Sci Rep* 2018;8(1):13630.
83. Omidbakhsh N, Sattar SA. Broad-spectrum microbicidal activity, toxicologic assessment, and materials compatibility of a new generation of accelerated hydrogen peroxide-based environmental surface disinfectant. *Am J Infect Control* 2006; 34(5):251–257.
84. Walsh LJ. Safety issues relating to the use of hydrogen peroxide in dentistry. *Aust Dent J* 2000;45(4):257–269.
85. Kraus FW, Perry WI, Nickerson JF. Salivary catalase and peroxidase values in normal subjects and in persons with periodontal disease. *Oral Surg Oral Med Oral Pathol* 1958;11(1): 95–102.
86. Cai J, Chen Y, Seth S, Furukawa S, Compans RW, Jones DP. Inhibition of influenza infection by glutathione. *Free Radic Biol Med* 2003;34(7):928–936.
87. Staal FJ, Roederer M, Herzenberg LA, Herzenberg LA. Intracellular thiols regulate activation of nuclear factor kappa B and transcription of human immunodeficiency virus. *Proc Natl Acad Sci USA* 1990;87(24):9943–9947.
88. Xiao J, Deng J, Lv L, et al. Hydrogen peroxide induce human cytomegalovirus replication through the activation of p38-MAPK signaling pathway. *Viruses* 2015;7(6):2816–2833.
89. Popkin DL, Zilka S, Dimaano M, et al. Cetylpyridinium chloride (CPC) exhibits potent, rapid activity against influenza viruses in vitro and in vivo. *Pathog Immun* 2017;2(2):252–269.
90. Kohn WG, Collins AS, Cleveland JL, et al. Guidelines for infection control in dental health-care settings–2003. *MMWR Recomm Rep* 2003;52(RR-17):1–61.
91. Feres M, Figueiredo LC, Faveri M, Stewart B, de Vizio W. The effectiveness of a preprocedural mouthrinse containing cetylpyridinium chloride in reducing bacteria in the dental office. *J Am Dent Assoc* 2010;141(4):415–422.
92. Retamal-Valdes B, Soares GM, Stewart B, et al. Effectiveness of a pre-procedural mouthwash in reducing bacteria in dental aerosols: randomized clinical trial. *Braz Oral Res* 2017;31:e21.
93. Meng L, Hua F, Bian Z. Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* 2020;99(5):481–487.
94. Ather A, Patel B, Ruparel NB, Diogenes A, Hargreaves KM. Coronavirus disease 19 (COVID-19): implications for clinical dental care. *J Endod* 2020;46(5):584–595.

95. Peng X, Xu X, Li Y, Cheng L, Zhou X, Ren B. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci* 2020;12(1):9.
96. Firquet S, Beaujard S, Lobert PE, et al. Survival of enveloped and non-enveloped viruses on inanimate surfaces. *Microbes Environ* 2015;30(2):140–144.
97. Jung MK, Callaci JJ, Lauing KL, et al. Alcohol exposure and mechanisms of tissue injury and repair. *Alcohol Clin Exp Res* 2011;35(3):392–399.
98. Simons K, Garoff H. The budding mechanisms of enveloped animal viruses. *J Gen Virol* 1980;50(1):1–21.
99. Sanchez EL, Lagunoff M. Viral activation of cellular metabolism. *Virology* 2015;479–480:609–618.
100. Wolff MH, Sattar SA, Adebunrin O, Tetro J. Environmental survival and microbicide inactivation of coronaviruses. In: A Schmidt, O Weber, MH Wolff, eds. *Coronaviruses with Special Emphasis on First Insights Concerning SARS*. Basel: Birkhäuser Basel; 2005:201–212.
101. Peiris JS, Yuen KY, Osterhaus AD, Stohr K. The severe acute respiratory syndrome. *N Engl J Med* 2003;349(25):2431–2441.
102. Slater JL, Huang, CH. Structure of biological membranes. In: P Yeagle, ed. *Structure of Biological Membranes*. Boca Raton, FL: CRC Press; 1992.
103. Slater JL, Huang CH. Interdigitated bilayer membranes. *Prog Lipid Res* 1988;27(4):325–359.
104. Eggers M. Infectious disease management and control with povidone iodine. *Infect Dis Ther* 2019;8(4):581–593.
105. van Meer G. Lipids of the Golgi membrane. *Trends Cell Biol* 1998;8(1):29–33.